

EPA Jacket 100-1411

Vol.2

Material Sent for Data Extraction

Reg. # 100-1411

Description: Accepted MFA-PEN-98-10

☒ Material(s) Sent to Data Extraction Contractors

☐ New Stamped Label Dated _____

☒ Notification Dated 3/26/2015

☒ New CSF(s) Dated 3/12/2015

☐ Other: _____

☒ Decision #: 503094

☐ Other Action/Comments: _____

File this coversheet and attached materials in the jacket. It must be well organized and clipped together, NOT STAPLED. Then give the jacket with the coversheet and materials to staff in the Information Services Center (ISC) (Room S-4900). If a jacket is full or only available as an image, please file materials in a new jacket and bring it down to the (ISC). For further information please call 703-605-0716.

Reviewer: Alganesh Debesai

Phone: 308-8353 Division: RD

Date: 4.22.15



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

April 21, 2015

Tammy Tyler, Ph. D.
Regulatory Product Manager
Syngenta Crop Protection, LLC.
P.O. Box 18300
Greensboro, NC 27419

Subject: Minor Formulation Amendment

Dear Dr. Tyler:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated 3/26/2015 for **EPA Registration 100-1411**. The Registration Division (RD) has conducted a review of two Confidential Statements of Formula (CSF) submitted with this request for its applicability under PRN 98-10 and finds that the change(s) requested falls within the scope of PRN 98-10. Therefore, revised Basic No. 1198/2 (A10324A) and new proposed Alternate No. 1199/2 (A10324A) CSFs dated 3/18/2015 **are acceptable**. A copy of each CSF has been added to the registration files for the subject product.

If you have any questions, please contact me via telephone at 703-308-8353 or e-mail (debesai.alganesh@epa.gov).

Sincerely,

Alganesh Debesai

Alganesh Debesai
Chemistry, Inerts and Toxicology Assessment Branch
Registration Division (7505P)
Office of Pesticide Programs

Please read instructions on reverse before completing form.

	United States Environmental Protection Agency Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number NOTIFICATION
---	--	--	--

Application for Pesticide - Section I

1. Company/Product Number 100-1411	2. EPA Product Manager Ms. Jessica Rogala	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Enfold™ Insecticide	PM# 7	
5. Name and Address of Applicant (Include ZIP Code) Syngenta Crop Protection, LLC P. O. Box 18300 Greensboro, NC 27419 <input type="checkbox"/> Check if this is a new address		6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____

Section - II

<input type="checkbox"/> Amendment - Explain below. <input type="checkbox"/> Resubmission in response to Agency letter dated _____ <input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____ <input type="checkbox"/> "Me Too" Application. <input type="checkbox"/> Other - Explain below.
--	---

Explanation: Use additional page(s) if necessary. (For Section I and Section II.).


Syngenta Crop Protection, LLC is submitting this notification to update the Enfold™ Insecticide (EPA Reg. No. 100-1411) Confidential Statement of Formula's (CSF's) listed suppliers.

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____	*Certification must be submitted	If "Yes" No. per Unit Packaging wgt. Container
If "Yes" No. per Unit Packaging wgt. Container		If "Yes" No. per Unit Packaging wgt. container			
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Other <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled					

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Tammy Tyler, Ph.D.	Title Regulatory Product Manager	Telephone No. (Include Area Code) 336-632-6055
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Data Application Received (Stamped)
2. Signature 	3. Title Regulatory Product Manager	
4. Typed Name Tammy Tyler, Ph.D.	5. Date March 26, 2015	

e-Submission



Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, NC 27419-8300
www.syngenta.com

FedEx

August 15, 2014

Document Processing Desk
Office of Pesticide Programs (7504P)
U.S. Environmental Protection Agency
Room S-4900, One Potomac Yard
2777 South Crystal Drive
Arlington, VA 22202-4501

Attention: Mr. Reuben Baris, Acting PM 7, Insecticide-Rodenticide Branch

<p>SUBJECT: ENFOLD EPA REG. NO. 100-1411 SUBMISSION OF FINAL PRINTED LABELING</p>
--

Dear Mr. Baris:

In accordance with the above subject matter, Syngenta Crop Protection, LLC is submitting one copy of final printed labeling for Enfold that reflects the new product labeling approved by EPA May 13, 2013.



Also enclosed is EPA Form 8570-1 indicating submission of final printed labeling. If you have any questions, please contact me at 336-632-7207. Thank you.

Sincerely,

A handwritten signature in cursive script that reads "Thomas J. Parshley".

Thomas J. Parshley
Senior Regulatory Product Manager

Enclosures

 United States Environmental Protection Agency Washington, DC 20460		<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number
Application for Pesticide - Section I			
1. Company/Product Number 100-1411		2. EPA Product Manager Reuben Baris	
4. Company/Product (Name) Enfold™		3. Proposed Classification <input type="checkbox"/> None <input checked="" type="checkbox"/> Restricted	
5. Name and Address of Applicant (Include ZIP Code) Syngenta Crop Protection, LLC P. O. Box 18300 Greensboro, NC 27419 <input type="checkbox"/> Check if this is a new address		6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	
Section - II			
<input type="checkbox"/> Amendment - Explain below. <input checked="" type="checkbox"/> Final printed labels in response to Agency letter dated <u>May 15, 2013</u> <input type="checkbox"/> Resubmission in response to Agency letter dated _____ <input type="checkbox"/> "Me Too" Application. <input type="checkbox"/> Notification - Explain below. <input type="checkbox"/> Other - Explain below.			
Explanation: Use additional page(s) if necessary. (For Section I and Section II.). Syngenta Crop Protection, LLC is submitting one copy of final printed labeling for Enfold that reflects the new product labeling approved by EPA May 13, 2013.			
Section - III			
1. Material This Product Will Be Packaged In:			
Child-Resistant Packaging <input type="checkbox"/> Yes* <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
*Certification must be submitted		If "Yes" Unit Packaging wgt. No. per Container _____	If "Yes" Unit Packaging wgt. No. per container _____
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 24 ounces	5. Location of Label Directions <input checked="" type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input checked="" type="checkbox"/> Other <u>Pressure Sensitive</u> <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			
Section - IV			
1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)			
Name Thomas J. Parshley		Title Senior Regulatory Product Manager	Telephone No. (Include Area Code) 336-632-7207
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.			6. Date Application Received (Stamped)
2. Signature 		3. Title Senior Regulatory Product Manager	
4. Typed Name Thomas J. Parshley		5. Date August 15, 2014	

RESTRICTED USE PESTICIDE

TOXIC TO FISH, MAMMALS, AND AQUATIC ORGANISMS

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION, AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

PULL HERE
TO OPEN

NOT REVIEWED
In Accordance with PR Notice 82-2
Based on Draft Labeling Dated
05/13/13



GROUP 6 INSECTICIDE

EnfoldTM

syngenta.

Insecticide

For control of listed lepidopterous larvae (worms/caterpillars) and suppression of leafminers and spider mites on outdoor-grown plants in commercial nursery production

Active Ingredient:

Emamectin benzoate (CAS No. 155569-91-8) 5.0%

Other Ingredients: 95.0%

Total: 100.0%

Enfold is a soluble granule containing 5% emamectin benzoate.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and directions for use in booklet.

EPA Reg. No. 100-1411 EPA Est. 67545-AZ-1

Product of China Formulated in USA

SCP 1411A-L1 0513
4040804

24 ounces
Net Weight



FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Have person sip glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.

continued...

FIRST AID (continued)

NOTE TO PHYSICIAN

Early signs of intoxication include dilation of pupils, muscular incoordination, and muscular tremors. Vomiting within one-half hour of exposure can minimize toxicity following accidental ingestion of the product; rapidly after exposure (<15 minutes), administer repeatedly medical charcoal in a large quantity of water or ipecac.

If toxicity from exposure has progressed to cause severe vomiting, the extent of resultant fluid and electrolyte imbalance should be gauged. Appropriate supportive parenteral fluid replacement therapy should be given, along with other required supportive measures (such as maintenance of blood pressure levels and proper respiratory functionality) as indicated by clinical signs, symptoms, and measurements.

In severe cases, observations should continue for at least several days until clinical condition is stable and normal. Since emamectin benzoate is believed to enhance GABA activity in animals, it is probably wise to avoid drugs that enhance GABA activity (barbiturates, benzodiazepines, valproic acid) in patients with potentially toxic emamectin benzoate exposure.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

HOT LINE NUMBER

For 24-Hour Medical Emergency Assistance (Human or Animal)
Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident)
Call

1-800-888-8372

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse. Avoid breathing dust or spray mist. Prolonged or frequently repeated exposure may cause allergic skin reactions in some individuals.

Personal Protective Equipment (PPE)

- **Ground and Handheld Application (except airblast sprayers):**

Mixers, loaders, and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber
- Shoes plus socks
- NIOSH-approved respirator with a dust-mist filter with MSAH/NIOSH approval number prefix TC-21 or any N, R, P, or HE filter.

Applicators must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber
- Shoes plus socks

- **Airblast Application:**

Mixers, loaders, and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber
- Shoes plus socks
- NIOSH-approved respirator with a dust-mist filter with MSAH/NIOSH approval number prefix TC-21 or any N, R, P, or HE filter.

continued...

PRECAUTIONARY STATEMENTS *(continued)*

Applicators using ENCLOSED CAB airblast sprayers

While inside the cab must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

When entering or leaving the cab must also wear:

- Chemical-resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber.

NOTE: Once inside the cab, applicator must remove gloves and store them in a chemical-resistant container such as a plastic bag.

- **Aerial application:**

Mixers, loaders, and other handlers must wear:

- Coveralls over long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber
- Dust/Mist-filtering respirator or a NIOSH-approved respirator with any N, R, P, or HE filter

Applicators (Enclosed Cockpit)

While inside the cockpit must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

When entering or leaving the cockpit must also wear:

- Chemical-resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber.

NOTE: Once inside the cockpit, applicator must remove gloves and store them in a chemical-resistant container such as a plastic bag.

Flaggers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170-240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations**Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothes immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

This pesticide is toxic to fish, birds, mammals, and aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment wash water or rinsate.

This product is highly toxic to bees if exposed to direct treatment or residues on blooming crops or other plants for up to 24 hours after application. To reduce potential exposure to pollinators that may be visiting the treatment area, do not apply this product or allow drift to blooming, pollen-shedding, or nectar-producing parts of plants during this time period.

Physical or Chemical Hazards

Do not use or store near heat or open flame.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Enfold must be used only in accordance with directions on this label or in separately published Syngenta supplemental labeling for this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber, or butyl rubber
- Shoes plus socks
- Protective eyewear

**FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL
MAY RESULT IN POOR INSECT CONTROL.**

GENERAL INFORMATION

Enfold is a selective insecticide for use on herbaceous and woody ornamental plants grown outdoors (in containers or in the ground) in commercial nursery production. Woody ornamentals include (but are not limited to) shrubs, non-bearing fruit and nut trees, Christmas trees, forest seedlings, and shade trees.

Enfold controls the larval stages (worms/caterpillars) of listed lepidopteran species and suppresses *Liriomyza* leafminer, Tetranychid mites and pear psylla. Enfold has contact activity, but is most efficacious when ingested by the pest. Shortly after exposure to Enfold, affected larvae are paralyzed, stop feeding, and subsequently die after 2-4 days.

- Apply Enfold to plant foliage when larvae first appear (immediately after egg hatch), but before populations reach damaging levels. Target Enfold applications at small (1/4 inch in length) larvae.
- Treatments must be made before larvae penetrate plant parts or before larvae begin webbing and sheltering.
- Thorough spray coverage is essential for optimum performance. Apply Enfold in sufficient water to ensure good coverage of all plant surfaces. The use of greater water volumes will generally result in better coverage, especially under adverse conditions (e.g., hot, dry) or when the plant canopy is dense.

Resistance Management

Enfold is a Group 6 insecticide (contains the active ingredient emamectin benzoate).

Because of the inherent risks of resistance development to any product, it is strongly advised that Enfold be used in a sound resistance management program. Treatment may not be effective against labeled pests if tolerant strains of insects or mites develop. When applying Enfold to plants that are hosts of labeled pests and these labeled pests have multiple generations per crop per year, use resistance management practices.

Resistance management practices may include, but are not limited to:

- Rotating Enfold with other products with different modes of action
- Avoiding treatment of successive pest generations with Enfold
- Using labeled rates at the specified spray intervals
- Using non-chemical alternatives such as beneficial arthropods
- Rotating susceptible to non-susceptible plants
- Using various cultural practices

For additional information regarding the implementation of these or other resistance management practices, consult your local agricultural advisor or company representative.

APPLICATION PROCEDURES

Application Prohibitions:

- **Chemigation:** Do not apply this product through any type of irrigation system.
- Open cab airblast application is prohibited; airblast applications must use an enclosed cab.
- **State Restriction:** Do not apply Enfold with aircraft in New York State.

Spray Equipment

Apply by ground, airblast sprayer or aircraft. Spray equipment configuration should be arranged to provide accurate, uniform, and thorough coverage of the target crop and minimize potential for spray drift. Use spray nozzles that provide medium to fine-sized droplets. To ensure accuracy, calibrate sprayer before each use. For spray equipment and calibration information, consult sprayer manufacturers and/or state recommendations. All ground and aerial application equipment must be properly maintained and calibrated using appropriate carriers.

Spray Volume

- Applications using sufficient water volume for thorough and uniform coverage of the target crop provide the most effective pest control.
- Avoid application when uniform coverage is not possible or if excessive spray drift or inversion is possible.

SPRAY DRIFT

NOTE: When states have more stringent regulations, they must be observed.

Spray Drift Precautions – Aircraft and Ground Application Equipment

Apply Enfold only when wind velocity favors on-target product deposition (approximately 3 to 10 mph).

- **Do not** apply with ground application equipment within 25 ft of or with aircraft within 150 ft of lakes, reservoirs, rivers, permanent streams, marshes, pot holes, natural ponds, estuaries, or commercial fish farm ponds.
- **Do not** cultivate within 25 ft of the aquatic area to allow growth of a vegetative filter strip.
- **Do not** allow this product to drift onto non-target areas. Drift may result in illegal residues or injury to non-target species. Risk of exposure to sensitive areas can be reduced by applying this product when the wind direction is away from the sensitive area.
- **Do not** apply when the weather conditions may cause drift:
 - Avoid application when the temperature is high and/or the humidity is low. These conditions increase the evaporation of spray droplets and the likelihood of drift to aquatic areas.
 - **Do not** apply when wind speed or wind gusts are greater than 10 mph.
 - **Do not** apply when wind speed is below 2 mph because wind direction will vary and there is a high potential for inversion.

Spray Drift Precautions (Aerial Application)

Responsibility

Avoiding spray drift at the application site is the responsibility of the applicator.

The interaction of many equipment- and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all of these factors when making application decisions.

Drift Management Requirements

The following drift management requirements must be followed to avoid off-target movement from aerial applications to non-target plants.

- **Outermost Nozzle Distance**

The distance of the outermost nozzles on the boom must not exceed $\frac{3}{4}$ the length of the wingspan or rotor.

- **Nozzle Direction**

Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

- **Maximum Wind Speed**

Do not apply when wind speed is greater than 10 mph.

- **Droplet Size**

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions. (See **Wind**, **Temperature and Humidity**, and **Temperature Inversions** sections.)

- **Controlling Droplet Size**

- Volume**

- Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

- Pressure**

- Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

- Number of Nozzles**

- Use the minimum number of nozzles that provide uniform coverage.

- Nozzle Orientation**

- Orienting nozzles so that the spray is released parallel to the air stream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.

- Nozzle Type**

- Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid-stream nozzles oriented straight back produce the largest droplets and the lowest drift.

- **Boom Length**

- For some use patterns, reducing the effective boom length to less than $\frac{3}{4}$ of the wingspan or rotor length may further reduce drift without reducing swath width.

- **Application Height**

- Applications should not be made at a height greater than 10 ft above the top of the target plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

- **Swath Adjustment**

When applications are made with a cross wind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind.

- **Wind**

Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application must be avoided below 2 mph due to variable wind direction and high inversion potential. **NOTE:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

- **Temperature and Humidity**

To compensate for evaporation when applying Enfold in low relative humidity, set up equipment to produce larger droplets. Evaporation of droplets is most severe when conditions are both hot and dry.

- **Temperature Inversions**

Enfold must not be applied during a temperature inversion because the potential for drift is high. Temperature inversions restrict vertical air mixing, and this causes small, suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds that are common during inversions. Temperature inversions are characterized by temperatures that increase with altitude and are common on nights with limited cloud cover and light to no wind. Inversions begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, the movement of smoke from a ground source or an aircraft smoke generator can also identify inversions. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates, indicates good vertical air mixing.

MIXING PROCEDURES

1. Thoroughly clean spray equipment before using this product.
2. Prepare no more spray mixture than is needed for the immediate application.
3. Keep product container tightly closed when not in use.
4. Agitate the spray solution before and during application.
5. Do not let the spray mixture stand overnight in the spray tank.
6. Flush the spray equipment thoroughly following each use and apply rinsate to a previously treated area.

Mixing Instructions: Enfold Alone

1. Add 1/3 of the required amount of water to the spray or mixing tank.
2. With the agitator running, add Enfold into the spray tank.
3. Continue agitation while adding the remainder of the water.
4. Begin application of the solution after Enfold has completely dispersed into the mix water.
5. Maintain agitation until all of the mixture has been applied.

Note: Do not use liquid fertilizer as a carrier for Enfold.

Enfold - Tank Mixtures

Compatibility

Enfold is compatible with most insecticide, fungicide, and foliar nutrient products. However, before tank mixing Enfold use a jar test, as described below, to test the physical compatibility of Enfold with tank mix partners.

1. Using a quart jar, add the proportionate amounts of the products to 1 qt of water. Add wettable powders and water-dispersible granular products first, then liquid flowables, and emulsifiable concentrates last.
2. After thoroughly mixing, let the mixture stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible.
3. If compatibility is acceptable, follow the instructions in **Mixing Instructions: Enfold Tank Mixtures**.

NOTE:

- If using Enfold in a tank mixture:
 - Do not mix with any product that prohibits such mixing.
 - Observe all directions for use, crop/sites, use rates, dilution ratios, precautions, and limitations that appear on the tank mix product label.
 - Do not exceed any labeled use rate.
 - Follow the most restrictive label precautions and limitations.
- Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are labeled.

Mixing Instructions: Enfold Tank Mixtures

1. Add 1/3 of the required amount of water to the mix tank.
2. Start the agitator running before adding any tank-mix partners.
3. When using Enfold in tank mixtures:
 - a. All products in water-soluble packaging should be added to the tank before any other tank-mix partner, including Enfold.
 - b. Allow the water-soluble packaging to completely dissolve and the product(s) to completely disperse before adding any other tank-mix partner to the tank.
 - c. Then add other tank-mix partners in this order: wettable powders, wettable granules (dry flowables), liquid flowables, liquids and emulsifiable concentrates. Always allow each tank mix partner to become fully dispersed before adding the next product.
4. Provide sufficient agitation while adding the remainder of the water.
5. Maintain agitation until all the mixture has been applied.

USE DIRECTIONS

- **Location Prohibition:** Do not use Enfold in greenhouses.
- **Number of Applications:** Do not apply more than 3 sequential applications of Enfold. Rotate to another insect control product with a different mode of action for at least two applications.
- **Adjuvant Recommendation:** Thorough spray coverage of plant foliage is essential for optimum control. To provide optimum coverage and insect control, the use of a penetrating type spray adjuvant such as horticultural spray oil (not a dormant oil) or a nonionic surfactant at the manufacturer's suggested rate is recommended. Do not use a sticker/binder type adjuvant or tank mix with products that contain a sticker/binder component in the formulation because this may reduce Enfold insect control.
- **Application following failure of another insecticide:** Do not apply Enfold following the failure of another product if the larvae are large (>1/4 inch long).

Plant Safety

NOTICE TO USER: Plant tolerance to Enfold has been found to be acceptable for many genera and species. Due to the large number of species and varieties of ornamentals and nursery plants, it is impossible to test every one for tolerance to Enfold. The professional user should determine if Enfold can be used safely prior to commercial use. In a small area, test the recommended rates on a small number of plants for phytotoxicity prior to widespread use.

Tank Mixture: The safety of all potential tank mixes on all plants may not have been tested. Before applying any tank mixture not specifically recommended on this label, the safety to the target plants should be confirmed.

Pest	Oz Product/A per Application	Instructions
Beet armyworm Cabbage looper Cabbage webworm Corn earworm Cross-striped cabbageworm Diamondback moth Fall armyworm Imported cabbageworm Southern armyworm Tobacco budworm Tobacco hornworm Tomato hornworm Tomato fruitworm Tomato pinworm Yellowstriped armyworm	2.4 – 4.8 oz/A	Apply when larvae are first observed. Application may be repeated at a 7- to 14-day interval to maintain control. Use 2.4 oz/A for low to moderate infestations and 4.8 oz/A for high infestations.
Alfalfa looper Apple pandemis Bagworm Bud moths: eyespotted tufted apple Cabbage looper Cankerworm species Codling moth Common winter moth European winter moth Fall webworm Filbertworm Fruitworms: cherry green species laconobia	3.2 – 4.8 oz/A	Apply when larvae are first observed. Application may be repeated at a 7- to 14-day interval to maintain control. Use 3.2 oz/A for low to moderate infestations and 4.8 oz/A for high infestations.

Pest	Oz Product/A per Application	Instructions
Genista caterpillar Gypsy moth Hickory shuckworm Leafminers: blister moth species tentiform species Leafrollers: filbert fruittree obliquebanded omnivorous redbanded variegated Lesser appleworm <i>Liriomyza</i> leafminers ¹ Mimosa webworm Navel orangeworm Peach twig borer Omnivorous leaftier Orange tortrix Oriental fruit moth Pear psylla ² Pecan bud moth Pecan casebearer species Pecan serpentine leafminer Redhumped caterpillar Soybean looper Spider mites ^{2,3} Spruce budworm Tent Caterpillars: Eastern Forest Walnut caterpillar	3.2 – 4.8 oz/A	Apply when larvae are first observed. Application may be repeated at a 7- to 14-day interval to maintain control. Use 3.2 oz/A for low to moderate infesta- tions and 4.8 oz/A for high infestations.

footnotes on next page...

- ¹ Enfold provides suppression of *Liriomyza trifolii*, *Liriomyza sativae*, and *Liriomyza hudriobrensis* populations. Suppression means either erratic control ranging from good to poor or consistent control at a level below that which is generally considered acceptable for commercial control.
- ² Enfold provides suppression. Suppression means either erratic control ranging from good to poor or consistent control at a level below that which is generally considered acceptable for commercial control.
- ³ Refers to phytophagous mites in the Acari subfamily Tetranychinae.

Enfold Mixing Aid and Spray Volume Dilution Chart

Ounces of Enfold Added to Specified Spray Solution Mix Volumes
Based on Listed Product Application Rate and Application Volume

Application Volume (Gallons per Acre)	Application Rate (Ounces of Product per Acre)	25 Gallons of Finished Spray	50 Gallons of Finished Spray	100 Gallons of Finished Spray
50	2.4	1.2	2.4	4.8
50	3.2	1.6	3.2	6.4
50	4.8	2.4	4.8	9.6
100	2.4	0.6	1.2	2.4
100	3.2	0.8	1.6	3.2
100	4.8	1.2	2.4	4.8
150	2.4	0.4	0.8	1.6
150	3.2	0.53	1.06	2.1
150	4.8	0.8	1.6	3.2
200	2.4	0.3	0.6	1.2
200	3.2	0.4	0.8	1.6
200	4.8	0.6	1.2	2.4

Application Volume (Gallons per Acre)	Application Rate (Ounces of Product per Acre)	25 Gallons of Finished Spray	50 Gallons of Finished Spray	100 Gallons of Finished Spray
250	2.4	0.24	0.48	0.96
250	3.2	0.32	0.64	1.28
250	4.8	0.48	0.96	1.92
300	2.4	0.2	0.4	0.8
300	3.2	0.26	0.53	1.06
300	4.8	0.4	0.8	1.6

1 ounce of Enfold = 9.5 teaspoons

Do not use household measuring utensils to measure Enfold.

Use Restrictions

- Allow a minimum of 7 days between applications.
- Do not apply more than 28.8 oz/A per year.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage

Store in a cool, dry area under lock and key. Post as a pesticide storage area. Always store pesticides in the original container. Store away from food, pet food, feed, seed, fertilizers, and veterinary supplies. Place liquid formulations on lower shelves and dry formulations above.

continued...

STORAGE AND DISPOSAL (continued)

Pesticide Disposal

Pesticide wastes may be hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Handling

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration.

Enfold™, the ALLIANCE FRAME
the SYNGENTA Logo
and the PURPOSE ICON
are Trademarks of a



Syngenta Group Company

©2014 Syngenta

For non-emergency (e.g., current product information) call
Syngenta Crop Protection at 1-800-334-9481.

Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina 27419-8300

SCP 1411A-L1 0513
4040804

RESTRICTED USE PESTICIDE

TOXIC TO FISH, MAMMALS, AND AQUATIC ORGANISMS

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION, AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

GROUP 6 INSECTICIDE



Enfold™

Insecticide

For control of listed lepidopterous larvae (worms/caterpillars) and suppression of leafminers and spider mites on outdoor-grown plants in commercial nursery production

Active Ingredient:

Emamectin benzoate
(CAS No. 155569-91-8) 5.0%

Other Ingredients: 95.0%

Total: 100.0%

Enfold Insecticide is a soluble granule containing 5% emamectin benzoate.

EPA Reg. No. 100-1411

EPA Est. 67545-AZ-1

Enfold™ and the SYNGENTA Logo are Trademarks of a Syngenta Group Company.

©2014 Syngenta

Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina 27419-8300

SCP 1411A-L1 0513
4040804

24 ounces
Net Weight

KEEP OUT OF REACH OF CHILDREN. CAUTION

See additional precautionary statements and directions for use in booklet.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to **FIRST AID** section in attached booklet for additional precautionary statements.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse. Avoid breathing dust or spray mist. Prolonged or frequently repeated exposure may cause allergic skin reactions in some individuals.

syngenta®



Environmental Hazards

This pesticide is toxic to fish, birds, mammals, and aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment wash water or rinsate.

This product is highly toxic to bees if exposed to direct treatment or residues on blooming crops or other plants for up to 24 hours after application. To reduce potential exposure to pollinators that may be visiting the treatment area, do not apply this product or allow drift to blooming, pollen-shedding, or nectar-producing parts of plants during this time period.

Physical or Chemical Hazards

Do not use or store near heat or open flame.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage

Store in a cool, dry area under lock and key. Post as a pesticide storage area. Always store pesticides in the original container. Store away from food, pet food, feed, seed, fertilizers, and veterinary supplies. Place liquid formulations on lower shelves and dry formulations above.

Pesticide Disposal

Pesticide wastes may be hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Handling

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration.

Chemigation: Do not apply this product through any type of irrigation system.

**SCP 1411A-L2 0513
4040800**

Material to be added to an e-Jacket/Jacket

Reg. # 100-1411

Decision # 449 308

Description: initial registration notice

☒ Material(s) Sent to Data Extraction Contractors:

☒ Stamped Label Dated 5/13/13

☐ Notification Dated _____

☐ New CSF(s) Dated _____

☐ Other: _____

☐ Other Action/Comments: _____

File this coversheet and attached materials in the jacket. It must be well organized and clipped together, NOT STAPLED. Then give the jacket with the coversheet and materials to staff in the Information Services Center (ISC) (Room S-4900). If a jacket is full or only available as an image, please file materials in a new jacket and bring it down to the (ISC). For further information please call 703-605-0716.

Reviewer: Tom Harris

Phone: 308-9423

Division: RD

Date: 5/13/13



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505C)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

100-1411

Date of Issuance:

MAY 13 2013

NOTICE OF PESTICIDE:

☒ Registration
☐ Reregistration
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

Enfold

Name and Address of Registrant (include ZIP Code):

Thomas J. Parshley
Syngenta Crop Protection, LLC
PO Box 18300
Greensboro, NC 27419

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/registration review of your product when the Agency requires all registrants of similar products to submit such data.

Signature of Approving Official:

Date:

MAY 13 2013

John Hebert, Product Manager 07
Insecticide-Rodenticide Branch, Registration Division (7505P)

EPA Form 8570-6

2. Make the following label changes before you release the product for shipment:

- Revise the EPA Registration Number to read, "EPA Reg. No 100-1411."

3. Submit one copy of the revised final printed label for the record before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF, dated 4/18/2011
- Alternate CSF #1, dated 4/18/2011

If you have any questions, please contact Tom Harris at 703-308-9423 or harris.thomas@epa.gov.

John Hebert
Product Manager 07
Insecticide-Rodenticide Branch
Registration Division (7505P)

Enclosure

(Booklet)

RESTRICTED USE PESTICIDE

TOXIC TO FISH, MAMMALS, AND AQUATIC ORGANISMS

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION, AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

Enfold™

Insecticide

GROUP 6 INSECTICIDE

For control of listed lepidopterous larvae (worms/caterpillars) and suppression of leafminers and spider mites on outdoor-grown plants in commercial nursery production

Active Ingredient:

Emamectin benzoate (CAS No. 155569-91-8)5.0%

Other Ingredients:95.0%

Total:100.0%

Enfold is a soluble granule containing 5% emamectin benzoate.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and directions for use in booklet.

EPA Reg. No. 100-XXXX

EPA Est. XXXXX

Product of XXXXX

Formulated in XXXXX

SCP

Net Weight

ACCEPTED

MAY 13 2013

**Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended, for the
pesticide registered under:**

EPA. Reg. No: 100-1411

FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Have person sip glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
<p align="center">NOTE TO PHYSICIAN</p> <p>Early signs of intoxication include dilation of pupils, muscular incoordination, and muscular tremors. Vomiting within one-half hour of exposure can minimize toxicity following accidental ingestion of the product; rapidly after exposure (< 15 minutes), administer repeatedly medical charcoal in a large quantity of water or ipecac.</p> <p>If toxicity from exposure has progressed to cause severe vomiting, the extent of resultant fluid and electrolyte imbalance should be gauged. Appropriate supportive parenteral fluid replacement therapy should be given, along with other required supportive measures (such as maintenance of blood pressure levels and proper respiratory functionality) as indicated by clinical signs, symptoms, and measurements.</p> <p>In severe cases, observations should continue for at least several days until clinical condition is stable and normal. Since emamectin benzoate is believed to enhance GABA activity in animals, it is probably wise to avoid drugs that enhance GABA activity (barbiturates, benzodiazepines, valproic acid) in patients with potentially toxic emamectin benzoate exposure.</p>	
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment.</p>	

HOT LINE NUMBER

For 24-Hour Medical Emergency Assistance (Human or Animal)
Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident)

Call

1-800-888-8372

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse. Avoid breathing dust or spray mist. Prolonged or frequently repeated exposure may cause allergic skin reactions in some individuals.

Personal Protective Equipment (PPE)

• **Ground and Handheld Application (except airblast sprayers):**

Mixers, loaders, and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber
- Shoes plus socks
- NIOSH-approved respirator with a dust-mist filter with MSAH/NIOSH approval number prefix TC-21 or any N, R, P, or HE filter.

Applicators must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber
- Shoes plus socks

• **Airblast Application:**

Mixers, loaders, and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber

- Shoes plus socks
- NIOSH-approved respirator with a dust-mist filter with MSAH/NIOSH approval number prefix TC-21 or any N, R, P, or HE filter.

Applicators using ENCLOSED CAB airblast sprayers

While inside the cab must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

When entering or leaving the cab must also wear:

- Chemical-resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber.

NOTE: Once inside the cab, applicator must remove gloves and store them in a chemical-resistant container such as a plastic bag.

- **Aerial application:**

Mixers, loaders, and other handlers must wear:

- Coveralls over long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber
- Dust/Mist-filtering respirator or a NIOSH-approved respirator with any N, R, P, or HE filter

Applicators (Enclosed Cockpit)

While inside the cockpit must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

When entering or leaving the cockpit must also wear:

- Chemical-resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber.

NOTE: Once inside the cockpit, applicator must remove gloves and store them in a chemical-resistant container such as a plastic bag.

Flaggers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170-240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothes immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

This pesticide is toxic to fish, birds, mammals, and aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment wash water or rinsate.

This product is highly toxic to bees if exposed to direct treatment or residues on blooming crops or other plants for up to 24 hours after application. To reduce potential exposure to pollinators that may be visiting the treatment area, do not apply this product or allow drift to blooming, pollen-shedding, or nectar-producing parts of plants during this time period.

Physical or Chemical Hazards

Do not use or store near heat or open flame.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Enfold must be used only in accordance with directions on this label or in separately published Syngenta supplemental labeling for this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber, or butyl rubber
- Shoes plus socks
- Protective eyewear

FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL.

GENERAL INFORMATION

Enfold is a selective insecticide for use on herbaceous and woody ornamental plants grown outdoors (in containers or in the ground) in commercial nursery production. Woody ornamentals include (but are not limited to) shrubs, non-bearing fruit and nut trees, Christmas trees, forest seedlings, and shade trees.

Enfold controls the larval stages (worms/caterpillars) of listed lepidopteran species and suppresses Liriomyza leafminer, Tetranychid mites and pear psylla. Enfold has contact activity, but is most efficacious when ingested by the pest. Shortly after exposure to Enfold, affected larvae are paralyzed, stop feeding, and subsequently die after 2-4 days.

- Apply Enfold to plant foliage when larvae first appear (immediately after egg hatch), but before populations reach damaging levels. Target Enfold applications at small (1/4 inch in length) larvae.
- Treatments must be made before larvae penetrate plant parts or before larvae begin webbing and sheltering.
- Thorough spray coverage is essential for optimum performance. Apply Enfold in sufficient water to ensure good coverage of all plant surfaces. The use of greater water volumes will generally result in better coverage, especially under adverse conditions (e.g., hot, dry) or when the plant canopy is dense.

Resistance Management

Enfold is a Group 6 insecticide (contains the active ingredient emamectin benzoate).

Because of the inherent risks of resistance development to any product, it is strongly advised that Enfold be used in a sound resistance management program. Treatment may not be effective against labeled pests if tolerant strains of insects or mites develop. When applying Enfold to plants that are hosts of labeled pests and these labeled pests have multiple generations per crop per year, use resistance management practices.

Resistance management practices may include, but are not limited to:

- Rotating Enfold with other products with different modes of action
- Avoiding treatment of successive pest generations with Enfold
- Using labeled rates at the specified spray intervals
- Using non-chemical alternatives such as beneficial arthropods
- Rotating susceptible to non-susceptible plants
- Using various cultural practices

For additional information regarding the implementation of these or other resistance management practices, consult your local agricultural advisor or company representative.

APPLICATION PROCEDURES

Application Prohibitions:

- **Chemigation:** Do not apply this product through any type of irrigation system.
- Open cab airblast application is prohibited; airblast applications must use an enclosed cab.
- **State Restriction:** Do not apply Enfold with aircraft in New York State.

Spray Equipment

Apply by ground, airblast sprayer or aircraft. Spray equipment configuration should be arranged to provide accurate, uniform, and thorough coverage of the target crop and minimize potential for spray drift. Use spray nozzles that provide medium to fine-sized droplets. To ensure accuracy, calibrate sprayer before each use. For spray equipment and calibration information, consult sprayer manufacturers and/or state recommendations. All ground and aerial application equipment must be properly maintained and calibrated using appropriate carriers.

Spray Volume

- Applications using sufficient water volume for thorough and uniform coverage of the target crop provide the most effective pest control.
- Avoid application when uniform coverage is not possible or if excessive spray drift or inversion is possible.

SPRAY DRIFT

NOTE: When states have more stringent regulations, they must be observed.

Spray Drift Precautions – Aircraft and Ground Application Equipment

Apply Enfold only when wind velocity favors on-target product deposition (approximately 3 to 10 mph).

- **Do not** apply with ground application equipment within 25 ft. of or with aircraft within 150 ft. of lakes, reservoirs, rivers, permanent streams, marshes, pot holes, natural ponds, estuaries, or commercial fish farm ponds.

- **Do not cultivate** within 25 ft. of the aquatic area to allow growth of a vegetative filter strip.
- **Do not allow** this product to drift onto non-target areas. Drift may result in illegal residues or injury to non-target species. Risk of exposure to sensitive areas can be reduced by applying this product when the wind direction is away from the sensitive area.
- **Do not apply** when the weather conditions may cause drift:
 - Avoid application when the temperature is high and/or the humidity is low. These conditions increase the evaporation of spray droplets and the likelihood of drift to aquatic areas.
 - **Do not apply** when wind speed or wind gusts are greater than 10 mph.
 - **Do not apply** when wind speed is below 2 mph because wind direction will vary and there is a high potential for inversion.

Spray Drift Precautions (Aerial Application)

Responsibility

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment- and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all of these factors when making application decisions.

Drift Management Requirements

The following drift management requirements must be followed to avoid off-target movement from aerial applications to non-target plants.

- **Outermost Nozzle Distance**
The distance of the outermost nozzles on the boom must not exceed $\frac{3}{4}$ the length of the wingspan or rotor.
- **Nozzle Direction**
Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.
- **Maximum Wind Speed**
Do not apply when wind speed is greater than 10 mph.
- **Droplet Size**
The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential

but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions. (See **Wind, Temperature and Humidity, and Temperature Inversions** sections.)

- **Controlling Droplet Size**

- Volume**

- Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

- Pressure**

- Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

- Number of Nozzles**

- Use the minimum number of nozzles that provide uniform coverage.

- Nozzle Orientation**

- Orienting nozzles so that the spray is released parallel to the air stream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.

- Nozzle Type**

- Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid-stream nozzles oriented straight back produce the largest droplets and the lowest drift.

- **Boom Length**

- For some use patterns, reducing the effective boom length to less than $\frac{3}{4}$ of the wingspan or rotor length may further reduce drift without reducing swath width.

- **Application Height**

- Applications should not be made at a height greater than 10 ft. above the top of the target plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

- **Swath Adjustment**

- When applications are made with a cross wind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind.

- **Wind**

- Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any

given speed. Application must be avoided below 2 mph due to variable wind direction and high inversion potential. **NOTE:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

- **Temperature and Humidity**

To compensate for evaporation when applying Enfold in low relative humidity, set up equipment to produce larger droplets. Evaporation of droplets is most severe when conditions are both hot and dry.

- **Temperature Inversions**

Enfold must not be applied during a temperature inversion because the potential for drift is high. Temperature inversions restrict vertical air mixing, and this causes small, suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds that are common during inversions. Temperature inversions are characterized by temperatures that increase with altitude and are common on nights with limited cloud cover and light to no wind. Inversions begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, the movement of smoke from a ground source or an aircraft smoke generator can also identify inversions. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates, indicates good vertical air mixing.

MIXING PROCEDURES

1. Thoroughly clean spray equipment before using this product.
2. Prepare no more spray mixture than is needed for the immediate application.
3. Keep product container tightly closed when not in use.
4. Agitate the spray solution before and during application.
5. Do not let the spray mixture stand overnight in the spray tank.
6. Flush the spray equipment thoroughly following each use and apply rinsate to a previously treated area.

Mixing Instructions: Enfold Alone

1. Add 1/3 of the required amount of water to the spray or mixing tank.
2. With the agitator running, add Enfold into the spray tank.
3. Continue agitation while adding the remainder of the water.
4. Begin application of the solution after Enfold has completely dispersed into the mix water.
5. Maintain agitation until all of the mixture has been applied.

Note: Do not use liquid fertilizer as a carrier for Enfold.

Enfold - Tank Mixtures

Compatibility

Enfold is compatible with most insecticide, fungicide, and foliar nutrient products. However, before tank mixing Enfold use a jar test, as described below, to test the physical compatibility of Enfold with tank mix partners.

1. Using a quart jar, add the proportionate amounts of the products to 1 qt. of water. Add wettable powders and water-dispersible granular products first, then liquid flowables, and emulsifiable concentrates last.
2. After thoroughly mixing, let the mixture stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible.
3. If compatibility is acceptable, follow the instructions in **Mixing Instructions: Enfold Tank Mixtures**.

NOTE:

- If using Enfold in a tank mixture:
 - Do not mix with any product that prohibits such mixing.
 - Observe all directions for use, crop/sites, use rates, dilution ratios, precautions, and limitations that appear on the tank mix product label.
 - Do not exceed any labeled use rate.
 - Follow the most restrictive label precautions and limitations.
- Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are labeled.

Mixing Instructions: Enfold Tank Mixtures

1. Add 1/3 of the required amount of water to the mix tank.
2. Start the agitator running before adding any tank-mix partners.
3. When using Enfold in tank mixtures:
 - a. All products in water-soluble packaging should be added to the tank before any other tank-mix partner, including Enfold.
 - b. Allow the water-soluble packaging to completely dissolve and the product(s) to completely disperse before adding any other tank-mix partner to the tank.
 - c. Then add other tank-mix partners in this order: wettable powders, wettable granules (dry flowables), liquid flowables, liquids and emulsifiable concentrates. Always allow each tank mix partner to become fully dispersed before adding the next product.
4. Provide sufficient agitation while adding the remainder of the water.
5. Maintain agitation until all the mixture has been applied.

USE DIRECTIONS

- **Location Prohibition:** Do not use Enfold in greenhouses.
- **Number of Applications:** Do not apply more than 3 sequential applications of Enfold. Rotate to another insect control product with a different mode of action for at least two applications.
- **Adjuvant Recommendation:** Thorough spray coverage of plant foliage is essential for optimum control. To provide optimum coverage and insect control, the use of a penetrating type spray adjuvant such as horticultural spray oil (not a dormant oil) or a nonionic surfactant at the manufacturer's suggested rate is recommended. Do not use a sticker/binder type adjuvant or tank mix with products that contain a sticker/binder component in the formulation because this may reduce Enfold insect control.
- **Application following failure of another insecticide:** Do not apply Enfold following the failure of another product if the larvae are large (>1/4 inch long).

Plant Safety

NOTICE TO USER: Plant tolerance to Enfold has been found to be acceptable for many genera and species. Due to the large number of species and varieties of ornamentals and nursery plants, it is impossible to test every one for tolerance to Enfold. The professional user should determine if Enfold can be used safely prior to commercial use. In a small area, test the recommended rates on a small number of plants for phytotoxicity prior to widespread use.

Tank Mixture: The safety of all potential tank mixes on all plants may not have been tested. Before applying any tank mixture not specifically recommended on this label, the safety to the target plants should be confirmed.

Pest	oz. Product/A per Application	Instructions
Beet armyworm Cabbage looper Cabbage webworm Corn earworm Cross-striped cabbageworm Diamondback moth Fall armyworm Imported cabbageworm Southern armyworm Tobacco budworm Tobacco hornworm Tomato hornworm Tomato fruitworm Tomato pinworm Yellowstriped armyworm	2.4-4.8 oz./A	Apply when larvae are first observed. Application may be repeated at a 7- to 14-day interval to maintain control. Use 2.4 oz./A for low to moderate infestations and 4.8 oz./A for high infestations.
Alfalfa looper Apple pandemis Bagworm Bud moths: eyespotted tufted apple Cabbage looper Cankerworm species Codling moth Common winter moth European winter moth Fall webworm Filbertworm Fruitworms: cherry green species laconobia Genista caterpillar Gypsy moth Hickory shuckworm Leafminers: blister moth species tentiform species Leafrollers: filbert fruittree obliquebanded omnivorous redbanded variegated Lesser appleworm <i>Liriomyza</i> leafminers ¹ Mimosa webworm Navel orangeworm Peach twig borer Omnivorous leaf-tier Orange tortrix Oriental fruit moth Pear psylla ² Pecan bud moth Pecan casebearer species	3.2-4.8 oz./A	Apply when larvae are first observed. Application may be repeated at a 7- to 14-day interval to maintain control. Use 3.2 oz./A for low to moderate infestations and 4.8 oz./A for high infestations.

Pecan serpentine leafminer		
Redhumped caterpillar		
Soybean looper		
Spider mites ^{2,3}		
Spruce budworm		
Tent Caterpillars:		
Eastern		
Forest		
Walnut caterpillar		

¹ Enfold provides suppression of *Liriomyza trifolii*, *Liriomyza sativae*, and *Liriomyza hudsonensis* populations. Suppression means either erratic control ranging from good to poor or consistent control at a level below that which is generally considered acceptable for commercial control.

² Enfold provides suppression. Suppression means either erratic control ranging from good to poor or consistent control at a level below that which is generally considered acceptable for commercial control.

³ Refers to phytophagous mites in the Acari subfamily Tetranychinae.

Enfold Mixing Aid and Spray Volume Dilution Chart

Ounces of Enfold Added to Specified Spray Solution Mix Volumes Based on Listed Product Application Rate and Application Volume

Application Volume (Gallons Per Acre)	Application Rate (Ounces of Product Per Acre)	25 Gallons of Finished Spray	50 Gallons of Finished Spray	100 Gallons of Finished Spray
50	2.4	1.2	2.4	4.8
50	3.2	1.6	3.2	6.4
50	4.8	2.4	4.8	9.6
100	2.4	0.6	1.2	2.4
100	3.2	0.8	1.6	3.2
100	4.8	1.2	2.4	4.8
150	2.4	0.4	0.8	1.6
150	3.2	0.53	1.06	2.1
150	4.8	0.8	1.6	3.2
200	2.4	0.3	0.6	1.2
200	3.2	0.4	0.8	1.6
200	4.8	0.6	1.2	2.4
250	2.4	0.24	0.48	0.96
250	3.2	0.32	0.64	1.28
250	4.8	0.48	0.96	1.92
300	2.4	0.2	0.4	0.8
300	3.2	0.26	0.53	1.06
300	4.8	0.4	0.8	1.6

1 ounce of Enfold = 9.5 teaspoons

Do not use household measuring utensils to measure Enfold.

Use Restrictions

- Allow a minimum of 7 days between applications.
- Do not apply more than 28.8 oz./A per year.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage

Store in a cool, dry area under lock and key. Post as a pesticide storage area. Always store pesticides in the original container. Store away from food, pet food, feed, seed, fertilizers, and veterinary supplies. Place liquid formulations on lower shelves and dry formulations above.

Pesticide Disposal

Pesticide wastes may be hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Handling

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration.

Enfold™, the ALLIANCE FRAME, the SYNGENTA Logo and the PURPOSE ICON are Trademarks of a Syngenta Group Company

©2012 Syngenta

For non-emergency (e.g., current product information) call
Syngenta Crop Protection at 1-800-334-9481.

Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina 27419-8300

SCP

(Non-detachable Container Label)

RESTRICTED USE PESTICIDE

TOXIC TO FISH, MAMMALS, AND AQUATIC ORGANISMS

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION, AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

Enfold™

Insecticide

GROUP 6 INSECTICIDE

For control of listed lepidopterous larvae (worms/caterpillars) and suppression of leafminers and spider mites on outdoor-grown plants in commercial nursery production

Active Ingredient:

Emamectin benzoate (CAS No. 155569-91-8).....5.0%

Other Ingredients:.....95.0%

Total:.....100.0%

Enfold Insecticide is a soluble granule containing 5% emamectin benzoate.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and directions for use in booklet.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

EPA Reg. No. 100-XXXX

EPA Est. XXXXX

Product of XXXXX

Formulated in XXXXX

SCP

Net Weight

Refer to **FIRST AID** section in attached booklet for additional precautionary statements.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse. Avoid breathing dust or spray mist. Prolonged or frequently repeated exposure may cause allergic skin reactions in some individuals.

Environmental Hazards

This pesticide is toxic to fish, birds, mammals, and aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment wash water or rinsate.

This product is highly toxic to bees if exposed to direct treatment or residues on blooming crops or other plants for up to 24 hours after application. To reduce potential exposure to pollinators that may be visiting the treatment area, do not apply this product or allow drift to blooming, pollen-shedding, or nectar-producing parts of plants during this period.

Physical or Chemical Hazards

Do not use or store near heat or open flame.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage

Store in a cool, dry area under lock and key. Post as a pesticide storage area. Always store pesticides in the original container. Store away from food, pet food, feed, seed, fertilizers, and veterinary supplies. Place liquid formulations on lower shelves and dry formulations above.

Pesticide Disposal

Pesticide wastes may be hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Handling

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration.

Chemigation: Do not apply this product through any type of irrigation system.

Enfold™ and the SYNGENTA Logo are Trademarks of a Syngenta Group Company.

©2013 Syngenta

Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina 27419-8300

SCP

Enfold (draft new product) – pl – 5/13/11
000100-XXXXX.20110516.ENFOLD_NEW-PRODUCT_MAY2011.pdf
Enfold (draft new product-version b) CLEAN – pl – 9/21/11
000100-XXXXX.20110921.ENFOLD_NEW-PRODUCT-B_SEP2011.pdf
Enfold (draft new product-c) CLEAN – pl – 3/22/12
000100-XXXXX.20120322.ENFOLD_NEW_C_MAR2012.pdf
Enfold (draft new product-d) CLEAN – pl – 3/27/12
000100-XXXXX.20120327.ENFOLD_NEW_D_MAR2012.pdf
Enfold (draft new product-e) –HILITE – pl – 4/25/13
000100-XRURR.20130425.ENFOLD_NEW_E_APR2013-HILITE.pdf
Enfold (draft new product-e) –CLEAN – pl – 4/25/13
000100-XRURR.20130425.ENFOLD_NEW_E_APR2013-CLEAN.pdf
Enfold (draft new product-f) – HILITE – pl – 4/30/13
000100-XRURR.20130430.ENFOLD_NEW_F_APR2013-HILITE.pdf
Enfold (draft new product-f) – CLEAN – pl – 4/30/13
000100-XRURR.20130430.ENFOLD_NEW_F_APR2013-CLEAN.pdf

